arXiv:1410.6919v1 [math.NT] 25 Oct 2014

THE RANK AND GENERATORS OF KIHARA'S ELLIPTIC CURVE WITH TORSION $\mathbb{Z}/4\mathbb{Z}$ OVER $\mathbb{Q}(t)$

ANDREJ DUJELLA, IVICA GUSIĆ AND PETRA TADIĆ

ABSTRACT. For the elliptic curve E over $\mathbb{Q}(t)$ found by Kihara, with torsion group $\mathbb{Z}/4\mathbb{Z}$ and rank ≥ 5, which is the current record for the rank of such curves, by using a suitable injective specialization, we determine exactly the rank and generators of $E(\mathbb{Q}(t))$.

1. Introduction

Let E be an elliptic curve over $\mathbb{Q}$. By the Mordell-Weil theorem, the group $E(\mathbb{Q})$ of rational points on E is a finitely generated abelian group. Hence, it is the product of the torsion group and $r \geq 0$ copies of the infinite cyclic group: $E(\mathbb{Q}) \cong E(\mathbb{Q})_{\text{tors}} \times \mathbb{Z}^r$. An important question in arithmetic geometry is to characterize which groups are possible as $E(\mathbb{Q})$ for an elliptic curve E over $\mathbb{Q}$. Thus, we may ask which torsion groups are possible, which ranks are possible, and, most ambitiously, which combinations of torsion and rank are possible. By Mazur's theorem, we know that $E(\mathbb{Q})_{\text{tors}}$ is one of the following 15 groups: $\mathbb{Z}/n\mathbb{Z}$ with $1 \leq n \leq 10$ or $n = 12$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2m\mathbb{Z}$ with $1 \leq m \leq 4$. On the other hand, it is not known which values of rank r are possible for elliptic curves over $\mathbb{Q}$. The folklore conjecture is that the rank can be arbitrary large, but it seems to be very hard to find examples with high rank. The current record is an example of an elliptic curve over $\mathbb{Q}$ with rank ≥ 28, found by Elkies in 2006. There is even a stronger conjecture that for any of 15 possible torsion groups T we have $B(T) = \infty$, where $B(T) = \sup\{\text{rank}\,(E(\mathbb{Q})) : \text{torsion group of } E \text{ over } \mathbb{Q} \text{ is } T\}$. The current records for ranks for each of the 15 possible torsion groups can be found on the web page [3].

Similar questions can be asked for elliptic curves over the field of rational functions $\mathbb{Q}(t)$. Again, we have 15 possible torsion groups, and we define $G(T) = \sup\{\text{rank}\, E(\mathbb{Q}(t)) : E(\mathbb{Q}(t))_{\text{tors}} \cong T\}$. In the next table, taken from

The authors were supported by the Croatian Science Foundation under the project no. 6422.

[0] *2010 Mathematics Subject Classification.* 11G05, 14H52.

Key words and phrases. Elliptic curve, specialization homomorphism, rank, torsion, generators, mwrank

[4], we give the current records for the rank of elliptic curves over $\mathbb{Q}(t)$ with prescribed torsion group.

T	$G(T) \geq$	Author(s)
0	18	Elkies (2006)
$\mathbb{Z}/2\mathbb{Z}$	11	Elkies (2009)
$\mathbb{Z}/3\mathbb{Z}$	7	Elkies (2007)
$\mathbb{Z}/4\mathbb{Z}$	5	Kihara (2004), Elkies (2007)
$\mathbb{Z}/5\mathbb{Z}$	3	Lecacheux (2001), Eroshkin (2009), MacLeod (2014)
$\mathbb{Z}/6\mathbb{Z}$	3	Lecacheux (2001), Kihara (2006), Eroshkin (2008), Woo (2008), Dujella & Peral (2012), MacLeod (2014)
$\mathbb{Z}/7\mathbb{Z}$	1	Kulesz (1998), Lecacheux (2003), Rabarison (2008), Harrache (2009), MacLeod (2014)
$\mathbb{Z}/8\mathbb{Z}$	2	Dujella & Peral (2012), MacLeod (2013)
$\mathbb{Z}/9\mathbb{Z}$	0	Kubert (1976)
$\mathbb{Z}/10\mathbb{Z}$	0	Kubert (1976)
$\mathbb{Z}/12\mathbb{Z}$	0	Kubert (1976)
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	7	Elkies (2007)
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$	4	Dujella & Peral (2012)
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$	2	Dujella & Peral (2012), MacLeod (2013)
$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$	0	Kubert (1976)

Note that in the previous table only the lower bounds for the rank are given. Indeed, it seems that only for the torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ the exact rank over $\mathbb{Q}(t)$ of the record curve can be found in literature. In fact, in [5], Dujella and Peral proved that the corresponding curve, obtained from the so called Diophantine triples, has rank equal to 4 and they provide the generators for the group. The proof uses the method introduced by Gusić and Tadić in [6] for an efficient search for injective specializations. In this paper, we will prove an analogous result for the curve with record rank over $\mathbb{Q}(t)$ with torsion group $\mathbb{Z}/4\mathbb{Z}$ found by Kihara [9] (see Theorem 2.1). Here we will use results from the recent paper [7], where the authors generalize and extend their method from [6]. In particular, by results of [7], now the method can be applied to curves with only one rational 2-torsion point.

Our main tool is [7, Theorem 1.3]. It deals with elliptic curves E given by $y^2 = x^3 + A(t)x^2 + B(t)x$, where $A, B \in \mathbb{Z}[t]$, with exactly one nontrivial 2-torsion point over $\mathbb{Q}(t)$. If $t_0 \in \mathbb{Q}$ satisfies the condition that for every nonconstant square-free divisor h of $B(t)$ or $A(t)^2 - 4B(t)$ in $\mathbb{Z}[t]$ the rational number $h(t_0)$ is not a square in $\mathbb{Q}$, then the specialized curve E_{t_0} is elliptic

and the specialization homomorphism at t_0 is injective. If additionally there exist $P_1, \ldots, P_r \in E(\mathbb{Q}(t))$ such that $P_1(t_0), \ldots, P_r(t_0)$ are the free generators of $E(t_0)(\mathbb{Q})$, then $E(\mathbb{Q}(t))$ and $E(t_0)(\mathbb{Q})$ have the same rank r, and $P_1, \ldots, P_r$ are the free generators of $E(\mathbb{Q}(t))$.

We can mention here that by the methods from [7], it is easy to show that the general families of curves with torsion $\mathbb{Z}/10\mathbb{Z}$, $\mathbb{Z}/12\mathbb{Z}$ and $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ given in Kubert's paper [10] all have rank 0 over $\mathbb{Q}(t)$, as expected from the corresponding entries in the above table (see Remark 3.1).

2. Kihara's curve with rank ≥ 5

In 2004, Kihara [9] constructed a curve over $\mathbb{Q}(t)$ with torsion group $\mathbb{Z}/4\mathbb{Z}$ and rank ≥ 5. This improved his previous result [8] with rank ≥ 4. We briefly describe Kihara's construction. The quartic curve H given by the equation

$$(x^2 - y^2)^2 + 2a(x^2 + y^2) + b = 0 \tag{2.1}$$

is considered. Forcing five points with coordinates of the form (r, s), (r, u), (s, p), (u, q), (p, m) to satisfy (2.1) leads to a system of certain quadratic Diophantine equations, for which a parametric solution is found. By the transformation $X = (a^2 - b)y^2/x^2$ and $Y = (a^2 - b)y(b + ax^2 + ay^2)/x^3$, we get from H the elliptic curve E with equation

$$Y^2 = X(X^2 + (2a^2 + 2b)X + (a^2 - b)^2). \tag{2.2}$$

We can write (2.2) in the form

$$Y^2 = X^3 + A(t)X^2 + B(t)X,$$

where

$$\begin{aligned}A(t) \;=\; & 26244t^{52} - 1454544t^{51} + 39753720t^{50} - 693511680t^{49} + 8373526532t^{48} - 71499362784t^{47} \\ & + 419498533496t^{46} - 1416310911728t^{45} - 901962029472t^{44} + 45707741530384t^{43} \\ & - 311918617874472t^{42} + 1112855854012288t^{41} - 445015546446924t^{40} - 2407470753825590 4t^{39} \\ & + 2017492091479359 60t^{38} - 1090449631523616720t^{37} + 4657677879944558620t^{36} \\ & - 1678972201074455004 8t^{35} + 5269149918548721841 6t^{34} - 1465292447259685158 40t^{33} \\ & + 3650829059462727442 64t^{32} - 8209368018922563460 48t^{31} + 1674263798355290924784t^{30} \\ & - 3107412066279198054752t^{29} + 5260774429646277796992t^{28} - 8138423149394923126752t^{27} \\ & + 1152424676987012782884 8t^{26} - 1497316243986572269056 0t^{25} + 1792653155650485238202 4t^{24} \\ & - 1993146445798516435385 6t^{23} + 2085143379750750528062 4t^{22} - 2091593799138116631952 0t^{21} \\ & + 2053473311547464286614 0t^{20} - 1997492217512773735220 8t^{19} + 1913799768894537755624 8t^{18} \\ & - 1763555588942563777510 4t^{17} + 1513669791417861862622 8t^{16} - 1172179695227266974422 4t^{15} \\ & + 7949459617497409864344t^{14} - 4573957484971788140208t^{13} + 2138285815612380734304t^{12} \\ & - 7488883271120272548 32t^{11} + 1527620052745984995 12t^{10} + 145618522077089516 80t^{9} \\ & - 26448842122489903356t^{8} + 1149341494890635248 0t^{7} - 299243384097172256 8t^{6} \\ & + 5118900432689670 24t^{5} - 57486161774987 812t^{4} + 4053242862 8328704t^{3} - 161507610671616t^{2} \\ & + 2684575014912t + 8707129344,\end{aligned}$$

$$\begin{aligned}B(t) \;=\; & 256t^2(t-1)^4(t+1)^4(t-3)^2(t-5)^2(3t-1)^2(t-2)^4(t^2+2)^2(t^2-2t+3)^2 \\ & \times (t^2+6t-1)^2(7t^2-18t+23)^2(3t^2-2t+7)^2(t^3+4t^2-5t+16)^2 \\ & \times (3t^4-17t^3+27t^2-43t+6)^2(2t^4-7t^3+9t^2-11t-5)^2(2t^4-17t^3+27t^2-25t+1)^2 \\ & \times (t^4-4t^3+6t^2-12t+1)^2(5t^4-17t^3+27t^2-79t+16)^2(t^4-28t^3+54t^2-92t+41)^2 \\ & \times (t^4-9t^3+15t^2-19t+4)^2.\end{aligned}$$

Kihara in [9] found five independent points $P_1, \ldots, P_5$ on this curve, corresponding to the five points on H mentioned above, showing that the rank of E over $\mathbb{Q}(t)$ is ≥ 5. The torsion subgroup is $\mathbb{Z}/4\mathbb{Z}$. Indeed, the point $T_1 = (a^2 - b, 2a(a^2 - b))$ on (2.2) is of order 4 since $2T_1 = (0, 0)$ and $4T_1 = \mathcal{O}$.

Our goal is to prove that the rank of E over $\mathbb{Q}(t)$ is exactly equal to 5 and to find the generators of $E(\mathbb{Q}(t))$. Computations with several specializations indicate that P_1, P_2, P_3, P_4, P_5 are not generators of $E(\mathbb{Q}(t))$. Indeed, from our results it will follow that they generate a subgroup of index 16 in $E(\mathbb{Q}(t))$.

In fact, it holds that $P_1 + P_i \in 2E(\mathbb{Q}(t))$ for $i = 2, 3, 4, 5$, i.e. there exist points W_2, W_3, W_4, W_5 of $E(\mathbb{Q}(t))$ such that $P_1 + P_i = 2W_i,\ i = 2, 3, 4, 5$. Since the torsion subgroup is $\mathbb{Z}/4\mathbb{Z}$, there are two choices for each W_i. We choose

one of them. The x-coordinates of these points are

$$\begin{aligned}
x(W_2) &= 4(t-3)(3t-1)(t^2+2)(t^4-4t^3+6t^2-12t+1)(3t^4-17t^3+27t^2-43t+6)\\
&\quad\times(t^4-9t^3+15t^2-19t+4)(7t^2-18t+23)(t^4-28t^3+54t^2-92t+41)(t-2)^2\\
&\quad\times(9t^{12}-110t^{11}+576t^{10}-2333t^9+7802t^8-19832t^7+39488t^6-57374t^5\\
&\quad+61421t^4-42914t^3+16488t^2-701t-216)^2(t+1)^4,\\
x(W_3) &= -16(t-1)^3(t+1)^3(t-3)(t^2+2)(t^2+6t-1)(t^3+4t^2-5t+16)\\
&\quad\times(t^4-9t^3+15t^2-19t+4)(2t^4-17t^3+27t^2-25t+1)(3t^4-17t^3+27t^2-43t+6)\\
&\quad\times(3t^2-2t+7)(t-2)^2t^2(3t-1)^2(t-5)^2(t^4-28t^3+54t^2-92t+41)^2\\
&\quad\times(2t^4-7t^3+9t^2-11t-5)^2,\\
x(W_4) &= 16t(t-1)^3(t+1)(3t-1)(t^2+2)(t^2+6t-1)(t^3+4t^2-5t+16)\\
&\quad\times(3t^4-17t^3+27t^2-43t+6)(t^4-9t^3+15t^2-19t+4)(2t^4-17t^3+27t^2-25t+1)\\
&\quad\times(3t^2-2t+7)(t-2)^2(t-5)^2(7t^2-18t+23)^2(t^4-4t^3+6t^2-12t+1)^2\\
&\quad\times(2t^4-7t^3+9t^2-11t-5)^2(3t^7-29t^6+51t^5-136t^4+175t^3-267t^2+179t-24)^2/\\
&\quad(8-55t-29t^2+103t^3-120t^4+69t^5-27t^6+3t^7)^2,\\
x(W_5) &= 16(t-1)^3(t-2)^3(t-3)^2(t^2+2)^2(3t^2-2t+7)^2(t^3+4t^2-5t+16)^2\\
&\quad\times(t^4-28t^3+54t^2-92t+41)(t^4-4t^3+6t^2-12t+1)(3t^4-17t^3+27t^2-43t+6)\\
&\quad\times(t^4-9t^3+15t^2-19t+4)(5t^4-17t^3+27t^2-79t+16)(2t^4-17t^3+27t^2-25t+1)\\
&\quad\times(t^2+6t-1)(t-5)(3t-1)(t+1)t.
\end{aligned}$$

We also give the x-coordinate of P_1:

$$\begin{aligned}
x(P_1) &= 16t(t-2)^2(t-3)(t-5)(3t-1)(t^2+2)(3t^2-2t+7)(7t^2-18t+23)\\
&\quad\times(t^2+6t-1)(t^3+4t^2-5t+16)(2t^4-17t^3+27t^2-25t+1)\\
&\quad\times(2t^4-7t^3+9t^2-11t-5)(3t^4-17t^3+27t^2-43t+6)(t^4-4t^3+6t^2-12t+1)\\
&\quad\times(t^2-2t+3)(5t^4-17t^3+27t^2-79t+16)(t^4-9t^3+15t^2-19t+4)\\
&\quad\times(t^4-28t^3+54t^2-92t+41)(3t^{13}-128t^{12}+1185t^{11}-5018t^{10}+13628t^9-27704t^8\\
&\quad+44162t^7-63956t^6+84827t^5-100976t^4+92061t^3-52802t^2+10662t-552)^2/\\
&\quad(12t^{11}-219t^{10}+1699t^9-7248t^8+21004t^7-45434t^6+72862t^5-90128t^4\\
&\quad+77496t^3-46283t^2+10095t-768)^2.
\end{aligned}$$

The points $P_1, W_2, \ldots, W_5$ are a natural guess for the generators, and we will show that this is indeed true by proving the following theorem in the next section.

Theorem 2.1. *The elliptic curve E over $\mathbb{Q}(t)$ has rank equal to 5 with free generators the points P_1, W_2, W_3, W_4, W_5 and the torsion group is $\mathbb{Z}/4\mathbb{Z}$.*

3. An injective specialization

As described in the introduction, we use [7, Theorem 1.3] to find rational numbers t_0 for which the specialization map at t_0 is injective. The condition is that for each nonconstant square-free divisor h of $B(t)$ or $A(t)^2-4B(t)$ in $\mathbb{Z}[t]$ the rational number $h(t_0)$ is not a square in $\mathbb{Q}$. The condition is easy to check, and we can find many rationals t_0 satisfying it. However, the coefficients of the curve E are polynomials with large degrees and coefficients. Thus, for the success of our approach, it is crucial to find suitable specialization t_0 of reasonably small height. Furthermore, we need a specialization for which the rank of E_{t_0} over $\mathbb{Q}$ is equal to 5, so it is reasonable to consider only specializations for which the the root number of E_{t_0} is -1 (conjecturally implying that the rank is odd).

We find that the specialization at $t_0=-\frac{11}{4}$ satisfies all requirements. It remains to compute the rank and generators of $E_{-11/4}$. For that purpose, we use the excellent program [2] of Cremona, which is included in the program package Sage [11]. By extending significantly the default precision (we use options `-p 800 -b 11`), we get the elliptic curve $E_{-11/4}$ over $\mathbb{Q}$, given by the equation

$$\begin{aligned}y^2 &= x^3+484371205173916954475505177386303655600428018856419825361x^2\\ &+396439739626376856220803587225531969795935240135441835411849245773275686438689322206293615089071659357199590400 00x,\end{aligned}$$

which is of rank 5 with five free generators $G_1,\dots,G_5$ and the generator of the torsion group T_0, given by their x-coordinates

$$\begin{aligned}x(T_0) &= -199107945503532544541442922607063711556495848199368164800\\ x(G_1) &= \frac{3512892979529333096658438292468696732246493248325984400000 0}{49}\\ x(G_2) &= -\frac{133158954443896697901506918015048543332676567688802003564886338908791003264350930560 00}{761597589972635906773076904 01}\\ x(G_3) &= -\frac{536936685248255028234168808334606137477987482678373004839681994240}{1416844881}\\ x(G_4) &= \frac{477616878094060797794543416082366648044181864476558959811339468000}{9223369}\\ x(G_5) &= \frac{72794607048562741902495446909805219644710035926500797206565704598920220473102284960 0}{13386309077372650899951050449}\end{aligned}$$

The rank 5 and generators are also confirmed in the most recent version V2.20-10 of Magma [1] (by the function `MordellWeilShaInformation` with option `SetClassGroupBounds("GRH")`). Now denote by $P_1^*,W_2^*,\dots,W_5^*$ the points obtained from $P_1,W_2,\dots,W_5$ after specialization $t\mapsto -\frac{11}{4}$. We easily get that

$$\begin{aligned}P_1^* &= T_0+G_5,\\ W_2^* &= T_0-G_3+G_4,\\ W_3^* &= T_0+G_1+G_2-G_3+G_4-G_5,\\ W_4^* &= T_0-G_2+G_3-G_4+G_5,\\ W_5^* &= T_0+G_4.\end{aligned}$$

Here all points are chosen up to sign of y-coordinates. It is easy to check that the matrix of this base change (modulo torsion) is of determinant ± 1 (the signs of the determinant depends on the choice of the signs of y-coordinates), so we see that $P_1^*, W_2^*, \ldots, W_5^*$ are free generators of the elliptic curve $E_{-11/4}$ over $\mathbb{Q}$. From the comments at the end of the introduction, we see that [7, Theorem 1.3] now implies that E has rank 5 over $\mathbb{Q}(t)$ and that P_1, W_2, W_3, W_4, W_5 are its free generators. Since E has a point of fourth order and the torsion group of $E_{-11/4}(\mathbb{Q})$ is $\mathbb{Z}/4\mathbb{Z}$, we conclude that the torsion group of $E(\mathbb{Q}(t))$ is also $\mathbb{Z}/4\mathbb{Z}$. □

Remark 3.1. Here we prove that the elliptic curves with torsion $\mathbb{Z}/10\mathbb{Z}$, $\mathbb{Z}/12\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$ and with rank ≥ 0 listed in Kubert's paper [10] have rank equal to 0 over $\mathbb{Q}(t)$.

For torsion group $\mathbb{Z}/10\mathbb{Z}$, the curve is given by the equation

$$y^2 = x(x^2 - (2t^2 - 2t + 1)(4t^4 - 12t^3 + 6t^2 + 2t - 1)x + 16t^5(t-1)^5(t^2 - 3t + 1)).$$

The specialization for $t_0 = 6$ satisfies the condition of [7, Theorem 1.3], and the specialized elliptic curve has rank 0, which proves our claim for that torsion group.

For torsion group $\mathbb{Z}/12\mathbb{Z}$, the curve is

$$y^2 = x(x^2 + (t^8 - 12t^6 - 48t^5 - 162t^4 - 480t^3 - 540t^2 - 624t - 183)x + 1024(t^2+3)^2(t+1)^6),$$

and the specialization which satisfies the condition of [7, Theorem 1.3] and has rank 0 is $t_0 = 11$.

Finally, for torsion group $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/8\mathbb{Z}$, we have the curve

$$y^2 = x(x + (t^2 - 1)^4)(x + 16t^4).$$

The specialization for $t_0 = 6$ satisfies the condition of [7, Theorem 1.1], and the specialized elliptic curve has rank 0, which proves our claim in that case.

References

[1] W. Bosma, J. Cannon, and C. Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput., 24 (1997), 235-265.

[2] J. E. Cremona, Algorithms for Modular Elliptic Curves, Cambridge Univ. Press, 1997.

[3] A. Dujella, *High rank elliptic curves with prescribed torsion*, `http://web.math.pmf.unizg.hr/ duje/tors/tors.html`

[4] A. Dujella, *Infinite families of elliptic curves with high rank and prescribed torsion*, `http://web.math.pmf.unizg.hr/ duje/tors/generic.html`

[5] A. Dujella and J. C. Peral, *High rank elliptic curves with torsion $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z}$ induced by Diophantine triples*, LMS J. Comput. Math. **17** (2014), 282–288.

[6] I. Gusić and P. Tadić, *A remark on the injectivity of the specialization homomorphism*, Glas. Mat. Ser. III **47** (2012), 265–275.

[7] I. Gusić and P. Tadić, *Injectivity of the specialization homomorphism of elliptic curves*, J. Number Theory, to appear.
[8] S. Kihara, *On the rank of elliptic curves with a rational point of order* 4, Proc. Japan Acad. Ser A Math. Sci. **80** (2004), 26–27.
[9] S. Kihara, *On the rank of elliptic curves with a rational point of order* 4*, II*, Proc. Japan Acad. Ser A Math. Sci. **80** (2004), 158–159.
[10] D. S. Kubert, *Universal bounds on the torsion of elliptic curves*, Proc. London Math. Soc. **33** (1976), 193–237.
[11] W. A. Stein et al., *Sage Mathematics Software (Version 6.0)*, The Sage Development Team, 2013, `http://www.sagemath.org`.

(Andrej Dujella) Department of Mathematics, University of Zagreb, Bijenička cesta 30, 10000 Zagreb, Croatia
E-mail address, Andrej Dujella: `duje@math.hr`

(Ivica Gusić) Faculty of Chemical Engin. and Techn., University of Zagreb, Marulićev trg 19, 10000 Zagreb, Croatia
E-mail address, Ivica Gusić: `igusic@fkit.hr`

(Petra Tadić) Martićeva 23, 10000 Zagreb, Croatia
E-mail address, Petra Tadić: `petra.tadic.zg@gmail.com`